arXiv:1506.03494v2 [math.PR] 24 Jun 2015

POISSON STATISTICS FOR MATRIX ENSEMBLES AT LARGE TEMPERATURE

FLORENT BENAYCH-GEORGES AND SANDRINE PÉCHÉ

Abstract. In this article, we consider β-ensembles, *i.e.* collections of particles with random positions on the real line having joint distribution

$$\frac{1}{Z_N(\beta)}|\Delta(\lambda)|^{\beta}e^{-\frac{N\beta}{4}\sum_{i=1}^{N}\lambda_i^2}\mathrm{d}\lambda,$$

in the regime where $\beta \to 0$ as $N \to \infty$. We briefly describe the global regime and then consider the local regime. In the case where $N\beta$ stays bounded, we prove that the local eigenvalue statistics, in the vicinity of any real number, are asymptotically to those of a Poisson point process. In the case where $N\beta \to \infty$, we prove a partial result in this direction.

1. Introduction

General β-ensembles are collections of particles with random positions on the real line with joint distribution

$$\frac{1}{Z_N(\beta)}|\Delta(\lambda)|^{\beta}e^{-\sum_{i=1}^{N}V(\lambda_i)}\mathrm{d}\lambda, \tag{1}$$

where $\mathrm{d}\lambda$ denotes the Lebesgue measure on $\mathbb{R}^N$, $\Delta(\lambda) := \prod_{1\le i<j\le N}(\lambda_j - \lambda_i)$, V is a potential with enough growth at infinity (like the Gaussian potential $V_G(x) = \frac{x^2}{2}$) and $Z_N(\beta)$ is a normalizing constant. Their study is initially motivated by some considerations from physics: the probability distribution can be viewed as the equilibrium measure of a one dimensional Coulomb gas, but they actually appear to be connected to a broad spectrum of mathematics and physics, such as random matrices, number theory, lattice gas theory, quantum mechanics and Selberg-type integrals. In the case where $\beta = 1$, 2 or 4, the probability measure of (1) is the joint distribution of the eigenvalues of a random $N \times N$ matrix M with density proportional to $e^{-\operatorname{Tr} V(M)}$ on the space of respectively real symmetric, complex Hermitian or quaternionic Hermitian matrices (see e.g. [6]). Besides, it was proved by Dumitriu and Edelman in [15] that when V is the Gaussian potential, for any $\beta > 0$, the probability measure of (1) is the joint distribution of the eigenvalues of the

---

Date: June 25, 2015.

2000 *Mathematics Subject Classification.* 15A52;60F05.

Key words and phrases. Random matrices, β-ensembles, Poisson point process.

random $N \times N$ tridiagonal matrix

$$H = \begin{pmatrix} g_1 & X_2 & & & \\ X_2 & g_2 & X_3 & & \\ & X_3 & \ddots & \ddots & \\ & & \ddots & \ddots & X_N \\ & & & X_N & g_N \end{pmatrix}, \qquad (2)$$

where the g_i's are some $N(0,1)$ variables and for all i, $X_i = \sqrt{Y_i}$, with Y_i distributed thanks to the $\Gamma((i-1)\beta/2)$ law, everything being independent. For general potential V, there is no random matrix representation.

In the classical cases ($\beta = 1, 2, 4$), the combinatorial structure and repulsive interaction has been well–understood for a long time via the theory of determinantal or Pfaffian processes (see e.g. [6] for references). The understanding of these asymptotic spectral statistics to the full class of parameters $\beta > 0$ has recently mobilized a lot of research. For general β, despite the lack of structure, some enormous progress has been accomplished recently. For fixed β, a few results are now known. First, it is known from [8] that the empirical eigenvalue distribution of the rescaled matrix $\frac{1}{\sqrt{N}}H$ converges weakly as $N \to \infty$ to a probability measure which is the semi-circle distribution in the case of Gaussian potential. The local eigenvalue statistics in the large N-limit are also quite well understood. In the Gaussian setting, at the edge of the spectrum, Ramírez, Rider and Virág have shown in [21] that the eigenvalues of $N^{1/6}(H - 2\sqrt{N}I)$ converge in distribution to those of the so-called stochastic Airy operator. In the bulk of the spectrum, the limiting spectral statistics are asymptotically defined in terms of the Sine-β process, which is again defined as the solution of a stochastic equation by Valkó and Virág in [26]. In particular the authors show that the Sine-β, which is translation invariant, has a geometric description in terms of the Brownian carousel, a deterministic function of the Brownian motion in the hyperbolic plane. Some advances on β-ensembles have also been made by Sosoe and Wang [22, 23] and Bao and Su in [7].
The question of universality for these statistics has now become an important matter of interest: some enormous progress has recently been accomplished by Bourgade, Erdös and Yau in [11, 12, 13, 10]. Therein the authors consider general β-ensembles (when the potential V is $\mathcal{C}^4$ and regular, or, in the first papers, convex and analytic). Assuming that the limiting spectral distribution (which depends on V) is supported on a single interval, they prove that the limiting eigenvalue statistics at the edge of the spectrum are given by the β-Tracy-Widom distribution. The universality in the bulk of the spectrum is also proved. Another point of view to tackle β-ensembles and in particular the quantitative aspect of the repulsion between eigenvalues has been developed in particular by Allez, Bouchaud and Guionnet in [3, 1]. They show in particular that when $\beta \leq 2$, β-ensembles can be seen as an N-dimensional process whose evolution is a mixing of that of N independent real Brownian motions and of that of a β-Dyson Brownian motion.

The scope of this article is to understand the spectral behavior, at microscopic scale, of β-ensembles in the case where $\beta \to 0$ and $N \to \infty$ (so that β depends on the dimension N). At macroscopic scale, such ensembles have been considered recently by [25] (see also the close model studied in [2]). Therein it is proved that when $\beta N \to c$ for some constant $c > 0$, the scaled empirical eigenvalue distribution of $\frac{1}{\sqrt{\beta}} H$ converges to the spectral measure of a deterministic Jacobi matrix, the density of which is explicit. When $\beta N \to \infty$, the limiting empirical eigenvalue distribution of converges to the semi-circle distribution. Local eigenvalue statistics have not been considered yet.

We here also consider the regime where $\beta \to 0$ and $N \to \infty$, but study the local eigenvalue statistics. In [18], Killip and Stoiciu have considered the same question for circular β-ensembles. More precisely they study CMV matrices (which are discrete one-dimensional Dirac-type operators) with random decaying coefficients. For rapidly decreasing coefficients, the eigenvalues have rigid spacing while in the case of slow decrease, the eigenvalues are distributed according to a Poisson process. More precisely, they prove that local eigenvalue statistics of β-circular ensembles when $\beta \to 0$ are in the large N limit those of a Poisson process.

For real-symmetric ensembles, the same question has recently been considered from a formal point of view. Indeed, in [4, 5], Allez and Dumaz considered the $\beta \to 0$ limit of the Sine-β process and of the β-Tracy-Widom distribution. The $\beta \to 0$ limit of the Sine-β is also considered by Leblé and Sefaty in [19]. The approach used by [19] is based on approximation theory while [4] use the diffusion representation of the Sine-β process to consider the limit $\beta \to 0$.

One would expect again to prove that when $\beta \to 0$ simultaneously to $N \to \infty$, the eigenvalues in the vicinity of a point u in the bulk of the spectrum exhibit Poisson statistics. In this text, we prove that this is true when $N\beta$ stays bounded as $N \to \infty$. In the case where $\beta \to 0$ but $N\beta \to \infty$, we have a partial result which formally implies the Poisson statistics in the bulk, but does not allow to get a complete proof.

In Figure 1, we compare this result with numerical simulations, giving a numerical evidence of the fact that the Poisson approximation works well (but gets less accurate as β grows).

Notation. For $u = u(N)$ and $v = v(N)$ some sequences,

$$u \ll v \iff u/v \underset{N\to\infty}{\longrightarrow} 0 \qquad ; \qquad u \sim v \iff u/v \underset{N\to\infty}{\longrightarrow} 1.$$

Acknowledgment. *The authors thank Alice Guionnet for her contribution to the proof and her useful suggestions for simplifying the arguments. We also thank Paul Bourgade for useful discussions.*



Figure 1. **Numerical comparison of the spacings of the λ_k's with the spacings in a Poisson point process:** histogram, in blue, of the debiazed spacings $(\lambda_{k+1}-\lambda_k)\sqrt{4-\lambda_k^2}$ for $0.05N \le k \le .95N$, with $\beta = N^c$, for several values of $c \in (-1, 0)$, compared to the density, in red, of the exponential law. The histogram of the spacings is drawn thanks to 10^3 independent realizations of the distribution of (4) for $N = 10^3$, (we ordered the λ_k's in an increasing way so that $\lambda_{k+1}-\lambda_k$ is actually the spacing between two consecutive particles). We see that the Poisson approximation works well, but gets less accurate as β grows (note however that all simulations of the last row were made with β almost equal to 1).

2. STATEMENT OF RESULTS

2.1. Presentation of the model.

For any $\alpha, \beta > 0$ and any $N \ge 1$, we define

$$Z_N(\alpha,\beta) := \int_{\lambda\in\mathbb{R}^N} \Delta(\lambda)^\beta \mathrm{e}^{-\frac{\alpha}{2}\sum_{i=1}^N \lambda_i^2}\mathrm{d}\lambda, \tag{3}$$

with $\Delta(\lambda) := \prod_{1\le i<j\le N} |\lambda_j - \lambda_i|$.

Let us now consider an exchangeable family $(\lambda_1,\ldots,\lambda_N)$ of random variables with joint law

$$P^{(N)}_{\alpha,\beta}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_N) := \frac{1}{Z_N(\alpha,\beta)}\Delta(\lambda)^\beta \mathrm{e}^{-\frac{\alpha}{2}\sum_{i=1}^N \lambda_i^2}\mathrm{d}\lambda_1\cdots\mathrm{d}\lambda_N \tag{4}$$

with $Z_N(\alpha,\beta)$ the normalization constant defined at (3).

2.2. **Tridiagonal model and relation between α and β.** Let

$$H=\frac{1}{\sqrt{\alpha}}\begin{pmatrix} g_1 & X_2 & & & \\ X_2 & g_2 & X_3 & & \\ & X_3 & \ddots & \ddots & \\ & & \ddots & \ddots & X_N \\ & & & X_N & g_N \end{pmatrix}, \tag{5}$$

where the g_i's are some $N(0,1)$ variables and for all i, $X_i=\sqrt{Y_i}$, with Y_i distributed thanks to the $\Gamma((i-1)\beta/2)$ law. We known, by [15] or Section 4.5 of [6], that $P^{(N)}_{\alpha,\beta}$ is the joint law of the eigenvalues of H.

Note that $\operatorname{Tr} H$ is centered and $\alpha\,\mathbb{E}\operatorname{Tr} H^2=N+\beta\frac{N(N-1)}{2}$, so that for

$$\alpha\sim 1+N\beta/2, \tag{6}$$

the empirical eigenvalue distribution of H has asymptotic first moments 0 and 1.

2.3. **Global and local regime for bounded $N\beta$.** The following proposition gives the limit of the empirical distribution of the λ_i's. The probability measure μ_γ in question here has been studied in [1, 25].

Proposition 2.1 (Global regime for bounded $N\beta$)**.** *Suppose that $N\beta\longrightarrow 2\gamma\geq 0$ as $N\to\infty$ and $\alpha\longrightarrow\gamma+1$.*

a) *Then under the law $P^{(N)}_{\alpha,\beta}$, the random probability measure*

$$\frac{1}{N}\sum_{i=1}^{N}\delta_{\lambda_i}$$

converges in probability to an even probability measure μ_γ on $\mathbb{R}$, depending only on γ, with moments m_k defined at (26), satisfying $m_2=1$.

b) *The measure μ_γ is absolutely continuous with respect to the Lebesgue measure on $\mathbb{R}$, with a density f_{μ_γ} that is bounded on any compact set, and satisfies, for all $x>0$,*

$$\mu_\gamma(\mathbb{R}\backslash[-x,x])\ \leq\ C_\gamma\frac{\mathrm{e}^{-\frac{\gamma+1}{2}x^2}}{x}$$

where C_γ is a constant depending only on γ.

c) *μ_γ depends continuously on $\gamma\geq 0$, is equal to $N(0,1)$ if and only if $\gamma=0$, and tends to the semicircle law with support $[-2,2]$ as $\gamma\to\infty$.*

d) *For each k, $\operatorname{Var}(N^{-1}\sum_{i=1}^{N}\lambda_i^k)=O(N^{-1})$.*

The following theorem gives the limit local behavior of the λ_i's.

Theorem 2.2 (Poisson limit for bounded $N\beta$)**.** *Suppose that* $N\beta \longrightarrow 2\gamma \geq 0$ *as* $N \to +\infty$ *and that* $\alpha \sim N\beta/2 + 1$*. Fix* $E \in \mathbb{R}$*. As* $N \to \infty$*, the point process*

$$\sum_{i=1}^{N} \delta_{N(\lambda_i - E)}$$

with $(\lambda_1, \ldots, \lambda_N) \sim P_{\alpha,\beta}^{(N)}$*, converges in distribution to the law of a Poisson point process with intensity* $\theta \, \mathrm{d}x$ *on* $\mathbb{R}$*, for*

$$\theta := \frac{(\gamma+1)^{\gamma+\frac{1}{2}}}{\sqrt{2\pi}\Gamma(\gamma+1)} \exp\left\{-\frac{\gamma+1}{2}E^2 + 2\gamma \int \log|E-x| \mathrm{d}\mu_\gamma(x)\right\}, \tag{7}$$

with μ_γ *is in Proposition 2.1.*

Remark 2.3. Note that the formula of θ given at (7) should agree with the density of μ_γ at E as given in [1, 25], but we were not able to prove it so far.

2.4. **Case where $\beta \gg N^{-1}$.** Using the fact that $N^2\beta \gg N$, one can easily adapt the proof of the following theorem from [6].

Theorem 2.4 (LDP for $\beta \gg N^{-1}$)**.** *Suppose that as* $N \to \infty$, $\alpha = \alpha(N)$ *and* $\beta = \beta(N)$ *are such that* $\alpha \sim N\beta/2$, $N\beta \gg 1$ *and* β *is bounded. Then for* $(\lambda_1, \ldots, \lambda_N)$ *distributed according to* $P_{\alpha,\beta}^{(N)}$*, the sequence of random probability measures* $L_N := \frac{1}{N}\sum_{i=1}^{N} \delta_{\lambda_i}$ *satisfies a LDP in the set of probability measure on* $\mathbb{R}$ *endowed with the weak topology with speed* $N^2\beta$ *and good rate function* I *defined by*

$$I(\mu) := \iint f(x,y)\mu^{\otimes 2}(\mathrm{d}x\mathrm{d}y) - \frac{3}{8}, \tag{8}$$

with $f : \mathbb{R}^2 \to \mathbb{R} \cup \{+\infty\}$ *is the function defined by*

$$f(x,y) = \frac{x^2+y^2}{8} - \frac{1}{2}\log|x-y|. \tag{9}$$

Moreover, the unique minimum of I *is achieved at the semicircle law*

$$\sigma := \frac{1}{2\pi}\sqrt{4-x^2}\mathbb{1}_{|x|\leq 2}\mathrm{d}x$$

and we have

$$\lim_{N\to\infty} \frac{1}{N^2\beta} \log Z_N(\alpha,\beta) = -\iint f(x,y)\sigma^{\otimes 2}(\mathrm{d}x\mathrm{d}y) = -\frac{3}{8}. \tag{10}$$

In the case where $N\beta \gg 1$, as far as the local regime is concerned, we only have the following partial result, inspired from Johansson's work in [17]. Below, we explain how formally, it allows to prove the convergence of local statistics to the ones of a Poisson point process and to identify its density.

Theorem 2.5. *Let $\beta = \beta(N)$ and $\alpha = \alpha(N)$ positive such that*

$$\frac{1}{N} \ll \beta \ll \frac{1}{\log N} \qquad ; \qquad N\beta - 2\alpha \ll 1. \tag{11}$$

Let $h : \mathbb{R} \to \mathbb{R}$ be a bounded function having 9 continuous bounded derivatives and $(\lambda_1, \ldots, \lambda_N)$ be distributed according to $P^{(N)}_{\alpha,\beta}$. Then as $N \to \infty$, we have

$$\log \int \mathrm{e}^{\beta \sum_{j=1}^N h(\lambda_j)} P^{(N)}_{\alpha,\beta}(\mathrm{d}\lambda_1, \ldots, \mathrm{d}\lambda_N) - N\beta \int h(t)\mathrm{d}\sigma(t) \longrightarrow \int h(t)\mathrm{d}\nu(t) \tag{12}$$

for

$$\nu := \frac{1}{2}(\delta_{-2} + \delta_2) - \mathbb{1}_{|x|\le 2} \frac{1}{\pi\sqrt{4-x^2}}\mathrm{d}x \tag{13}$$

Remark 2.6. The measure ν of (13) is a classic correction to the semi-circle law (see e.g. [17, Rem. 2.5] or, more recently, [16]).

Let us now explain how, on the formal level, Theorem 2.5 gives, for any $E \in (-2, 2)$, the convergence of the point process

$$\sum_{i=1}^N \delta_{N(\lambda_i - E)}$$

to a Poisson point process with density $\theta \mathrm{d}x$ on $\mathbb{R}$, for

$$\theta := \frac{\sqrt{4-x^2}}{2\pi}, \tag{14}$$

The first thing one has to notice is that

$$\theta = \frac{1}{2\pi} \exp \int \log |E - t| \mathrm{d}\nu(t) \tag{15}$$

for ν being as in (13) (the proof goes along the same lines as [8, Lem. 2.7]).

To prove it rigorously, we would need to prove that for

$$\widetilde{R}^{(N)}_k(x_1, \ldots, x_k) := \int \mathrm{e}^{\beta \sum_{i=1}^k \sum_{j=1}^{N-k} \log|E + \frac{x_i}{N} - \lambda_j|} P^{(N-k)}_{\alpha,\beta}(\mathrm{d}\lambda_1, \ldots, \mathrm{d}\lambda_{N-k}), \tag{16}$$

for any $x_1, \ldots, x_k \in \mathbb{R}$,

$$\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \mathrm{e}^{-\frac{\alpha}{2}\sum_i (E + \frac{x_i}{N})^2} \widetilde{R}^{(N)}_k(x_1, \ldots, x_k) \longrightarrow \theta^k \tag{17}$$

and that we have an upper bound of the type of (64).

First, it can be proved (see Section 5.1) that as $N \to \infty$, for any fixed k,

$$\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \sim \left(\frac{\mathrm{e}^{N\beta/2}}{2\pi}\right)^k. \tag{18}$$

Moreover, there is a universal positive constant M independent of N such that uniformly on N, k,

$$\mathbb{1}_{1\leq k\leq N}\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \leq M^k\left(\frac{\mathrm{e}^{N\beta/2}}{2\pi}\right)^k. \tag{19}$$

Theorem 2.5 can be rewritten as follows: for each fixed h as in the theorem,

$$\mathbb{E}_{P^{(N)}_{\alpha,\beta}}\mathrm{e}^{\beta\sum_{j=1}^N h(\lambda_j)} = \exp\left\{N\beta\int h(t)\mathrm{d}\sigma(t) + \int h(t)\mathrm{d}\nu(t) + \varepsilon_N(h)\right\}, \tag{20}$$

with $\varepsilon_N(h) \ll 1$. By (20), cutting on the right thanks to Lemma 5.1 and making as if the function $h_N : \lambda \mapsto \sum_{i=1}^k \log|E + \frac{x_i}{N} - \lambda|$ were $\mathcal{C}^9$ (and close enough to the function $h : \lambda \mapsto k\log|E-\lambda|$), we should have

$$\begin{aligned}
\widetilde{R}_k^{(N)}(x_1,\ldots,x_k) &= \int \mathrm{e}^{N\beta\sum_{i=1}^{N-k}h_N(\lambda_j)}P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_{N-k})\\
&= \exp\left\{N\beta\int h_N(t)\mathrm{d}\sigma(t) + \int h_N(t)\mathrm{d}\nu(t) + \varepsilon_N(h_N)\right\}\\
&\approx \exp\left\{N\beta\int h(t)\mathrm{d}\sigma(t) + \int h(t)\mathrm{d}\nu(t) + \varepsilon_N(h)\right\}
\end{aligned}$$

for ν as in (13). But by [8] p. 529, we know that for any $E \in (-2,2)$,

$$\int k^{-1}h(t)\mathrm{d}\sigma(t) = \int \log|E-t|\mathrm{d}\sigma(t) = \frac{E^2}{4} - \frac{1}{2},$$

so that we should have

$$\widetilde{R}_k^{(N)}(x_1,\ldots,x_k) \approx \exp\left\{N\beta\left(\frac{kE^2}{4}-\frac{k}{2}\right) + \int h(t)\mathrm{d}\nu(t) + \varepsilon_N(h)\right\}. \tag{21}$$

Besides, by (18), we have

$$\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \sim \left(\frac{\mathrm{e}^{N\beta/2}}{2\pi}\right)^k. \tag{22}$$

Puting together (21), (22) and the fact that $\alpha \sim \frac{N\beta}{2}$, we should have

$$\begin{aligned}
\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)}\mathrm{e}^{-\frac{\alpha}{2}\sum_i(E+\frac{x_i}{N})^2}\widetilde{R}_k^{(N)}(x_1,\ldots,x_k) &\approx \left(\frac{\mathrm{e}^{N\beta/2}}{2\pi}\right)^k\mathrm{e}^{-\frac{kN\beta}{4}E^2}\mathrm{e}^{N\beta\left(\frac{kE^2}{4}-\frac{k}{2}\right)+\int h(t)\mathrm{d}\nu(t)}\\
&= \theta^k
\end{aligned} \tag{23}$$

with θ as in (15).

3. Proof of Proposition 2.1

Let H be as in (5). We shall prove that for any k, $N^{-1}\,\mathbb{E}\,\mathrm{Tr}\,H^k$ tends to m_k, that the m_k's satisfy Carleman's criterion and that d) holds. By Skorohod's representation theorem (see e.g. [14, Th. 2.3.2]) and a diagonal extraction, it will imply a). Part c) will be clear from the proof. Note first that if Y is a $\Gamma(t)$-distributed variable, then for all $k \geq 0$,

$$\mathbb{E}\,Y^k = \frac{\Gamma(t+k)}{\Gamma(t)}. \tag{24}$$

Let us fix $k \geq 0$, $u \in [0,1]$, let $i = i(N)$ be such that $i/N \longrightarrow u$ and compute $\mathbb{E}(H^k)_{ii}$. We have

$$(H^k)_{ii} = \sum_{\varepsilon} H_{\varepsilon(0),\varepsilon(1)} \cdot \cdots \cdot H_{\varepsilon(k-1),\varepsilon(k)},$$

where the sum is taken over paths $\varepsilon : \{0,\ldots,k\} \to \{1,\,\ldots,n\}$ such that

- $\varepsilon(0) = \varepsilon(k) = i$,
- for all $\ell = 1,\ldots,k$, $\varepsilon(\ell) - \varepsilon(\ell-1) = -1, 0$ or 1, in which case we say that ℓ belongs respectively to $\mathrm{D}(\varepsilon)$, $\mathrm{F}(\varepsilon)$ or $\mathrm{U}(\varepsilon)$.

Note first that for such a path ε,

$$\#\,\mathrm{D}(\varepsilon) = \#\,\mathrm{U}(\varepsilon), \qquad \#\,\mathrm{D}(\varepsilon) + \#\,\mathrm{F}(\varepsilon) + \#\,\mathrm{U}(\varepsilon) = k. \tag{25}$$

For any $j \in \{1,\ldots,N\}$, we introduce

$$\begin{aligned} \mathrm{F}_j(\varepsilon) &:= \{\ell \in \{1,\ldots,k\}\,;\, \varepsilon(\ell-1) = \varepsilon(\ell) = j\} \\ \mathrm{U}_j(\varepsilon) &:= \{\ell \in \mathrm{U}(\varepsilon)\,;\, \varepsilon(\ell) = j\} \end{aligned}$$

Then one can easily see, using (24), that

$$\begin{aligned} \alpha^{k/2}\,\mathbb{E}\,H_{\varepsilon(0),\varepsilon(1)} \cdot \cdots \cdot H_{\varepsilon(k-1),\varepsilon(k)} &= \prod_j (\mathbb{E}\,g^{\#\,\mathrm{F}_j(\varepsilon)}\,\mathbb{E}\,Y_j^{\#\,\mathrm{U}_j(\varepsilon)}) \\ &= \mathbb{1}_{\forall j,\ \#\,\mathrm{F}_j(\varepsilon) \text{ is even}} \times \\ &\quad \prod_j \left(\#\,\mathrm{F}_j(\varepsilon)!! \times \frac{\Gamma\left(\frac{(j-1)\beta}{2} + \#\,\mathrm{U}_j(\varepsilon)\right)}{\Gamma(\frac{(j-1)\beta}{2})} \right) \\ &\xrightarrow[N\to\infty]{} \mathbb{1}_{\forall j,\ \#\,\mathrm{F}_j(\varepsilon) \text{ is even}} \times \\ &\quad \prod_j (\#\,\mathrm{F}_j(\varepsilon)!! \times \gamma u(\gamma u+1)\cdots(\gamma u + \#\,\mathrm{U}_j(\varepsilon) - 1)) \end{aligned}$$

By the dominated convergence theorem, it follows that $N^{-1}\,\mathbb{E}\operatorname{Tr} H^k$ converges to

$$m_k := \sum_{\varepsilon} (\gamma+1)^{-k/2} \int_{u=0}^{1} \prod_j \{\#\,\mathrm{F}_j(\varepsilon)!! \times \gamma u(\gamma u+1)\cdots(\gamma u + \#\,\mathrm{U}_j(\varepsilon)-1)\}\,\mathrm{d}u, \tag{26}$$

where the sum runs over paths $\varepsilon : \{0,\ldots,k\} \to \mathbb{Z}$ whose steps are in $\{-1,0,1\}$, such that $\varepsilon(0)=\varepsilon(k)=0$ and for all k, $\#\,\mathrm{F}_j(\varepsilon)$ is even.

Note that $\sum_j \#\,\mathrm{F}_j(\varepsilon) = \#\,\mathrm{F}(\varepsilon)$, whose parity is the one of k by (25), so that when k is odd, $\mathbb{E}(H^k)_{ii}=0$. Using (25) again, we see that when k is even, for any ε, for any u,

$$\prod_j (\#\,\mathrm{F}_j(\varepsilon)!! \times \gamma u(\gamma u+1)\cdots(\gamma u + \#\,\mathrm{U}_j(\varepsilon)-1)) \le \max\{1,\gamma\}^k k!!,$$

so that the m_k's satisfy Carleman's criterion. It follows that the m_k's are the moments of a unique measure μ_γ which depends continuously on γ. Besides, d) follows from the fact that $\operatorname{Cov}((H^k)_{ii},(H^k)_{jj})=0$ as soon as $|j-i|>2k$.

If $\gamma=0$, then the only way for the term associated to ε in (26) to be non zero is that k is even and ε is the constant path equal to i. This proves that $\mu_0 = N(0,1)$. The reciprocal is obvious, as the fact that μ_γ tends to the semicircle law when $\gamma\to\infty$ (using the formula of the moments of the semicircle law in terms of Dyck paths, as in [6] or [20]).

To prove the first part of b), we use Lemma 4.2 below. For any $a\in\mathbb{R}$ and $\varepsilon\in(0,1]$, we have

$$\mu((a-\varepsilon,a+\varepsilon)) \le \liminf_{N\to\infty} P^{(N)}_{\alpha,\beta}(|\lambda_1 - a| < \varepsilon) \le C\varepsilon \mathrm{e}^{\frac{\gamma a^2}{4}}.$$

The second part of b) is a direct consequence of Lemma 4.3 below.

4. Proof of Theorem 2.2

4.1. **Correlation functions.** To prove the theorem, according to Proposition 6.1, we introduce the correlation functions of the point process $\sum_{i=1}^N \delta_{N(\lambda_i-E)}$, given by the formulas

$$R_k^{(N)}(x_1,\ldots,x_k) \;:=\; \frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)}\,\frac{N^{-k}N!}{(N-k)!}\,\frac{\Delta(x_1,\ldots,x_k)^\beta}{N^{\frac{\beta k(k-1)}{2}}}\mathrm{e}^{-\frac{\alpha}{2}\sum_i (E+\frac{x_i}{N})^2}\,\widetilde{R}_k^{(N)}(x_1,\ldots,x_k) \tag{27}$$

with

$$\widetilde{R}_k^{(N)}(x_1,\ldots,x_k) \;:=\; \int \mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log|E+\frac{x_i}{N}-\lambda_j|}\,P^{(N-k)}_{\alpha,\beta}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_{N-k}) \tag{28}$$

First of all, we know that

$$\mathbb{1}_{k\le N}\frac{N^{-k}N!}{(N-k)!} \le 1 \tag{29}$$

and that as $N \to \infty$, for each fixed k,

$$\frac{N^{-k} N!}{(N-k)!} \underset{N\to\infty}{\longrightarrow} 1. \tag{30}$$

Besides, for any $M > 0$, for any $k \geq 1$ and any $x_1, \dots, x_k \in [-M, M]$, we have

$$\mathbb{1}_{k\leq N} \frac{\Delta(x_1, \dots, x_k)^\beta}{N^{\frac{\beta k(k-1)}{2}}} \leq \left(\frac{(2M)^\beta}{N^{\frac{N\beta}{2}}} \right)^k \tag{31}$$

and as soon as $\beta \ll (\log N)^{-1}$, for any fixed k and any fixed $x_1, \dots x_k$,

$$\frac{\Delta(x_1, \dots, x_k)^\beta}{N^{\frac{\beta k(k-1)}{2}}} \underset{N\to\infty}{\longrightarrow} 1. \tag{32}$$

4.2. **Partition functions.** We know, by [6, Cor. 2.5.9], that

$$Z_N(\alpha, \beta) \;=\; \alpha^{-\left(\frac{N(N-1)}{4}\beta + \frac{N}{2}\right)} (2\pi)^{N/2} \prod_{j=1}^{N} \frac{\Gamma(1 + j\beta/2)}{\Gamma(1 + \beta/2)} \tag{33}$$

Hence

$$\begin{aligned}
\mathbb{1}_{k\leq N} \frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} &= \alpha^{\frac{k}{2}} \alpha^{\frac{\beta}{4}(N(N-1)-(N-k)(N-k-1))} \prod_{j=N-k+1}^{N} (2\pi)^{-1/2} \frac{\Gamma(1+\frac{\beta}{2})}{\Gamma(1+\frac{j\beta}{2})} \\
&= \alpha^{\frac{k}{2}} \alpha^{\frac{\beta}{4}(2kN-k(k+1))} \prod_{j=N-k+1}^{N} (2\pi)^{-1/2} \frac{\Gamma(1+\frac{\beta}{2})}{\Gamma(1+\frac{j\beta}{2})}
\end{aligned}$$

By hypothesis, $N\beta$ is bounded and so is α. Let $C \geq 1$ be such that $N\beta + \alpha \leq C$. Then we have, uniformly in k,

$$\mathbb{1}_{k\leq N} \frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \leq \left(\frac{C^{\frac{1+C}{2}}}{\sqrt{2\pi}} \frac{\max_{[1,C]} \Gamma}{\min_{[1,C]} \Gamma} \right)^k \tag{34}$$

Besides, as $N\beta \longrightarrow 2\gamma \geq 0$, for each fixed k,

$$\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \longrightarrow \left(\frac{(\gamma+1)^{\gamma+\frac{1}{2}}}{\sqrt{2\pi}\Gamma(\gamma+1)} \right)^k. \tag{35}$$

4.3. **Uniform upper-bound on the correlation functions.**

Lemma 4.1. *Let $\mathcal{K}$ be a compact subset of $\mathbb{R}$. There is a constant C depending only on $\mathcal{K}$ and on the upper bounds on the sequences $N\beta, \alpha$ such that for all k, N and all $x_1, \dots, x_k \in \mathcal{K}$, we have*

$$\mathbb{1}_{k\leq N} \widetilde{R}_k^{(N)}(x_1, \dots, x_k) \leq C^k \mathrm{e}^{\frac{1}{4}\sum_{i=1}^k x_i^2}.$$

Proof. Note that by (36) and (37), for each $i \in \{1, \ldots k\}$ and $j \in \{1, \ldots, N-k\}$, we have

$$|E + \frac{x_i}{N} - \lambda_j|^\beta \leq (|E + \frac{x_i}{N}| + |\lambda_j|)^\beta \leq 2^\beta \exp\{\beta \frac{(E + \frac{x_i}{N})^2 + \lambda_j^2}{8}\}.$$

Hence for C a constant (that might change from line to line) as in the statement of the lemma,

$$\begin{aligned} \widetilde{R}_k^{(N)}(x_1, \ldots, x_k) &\leq 2^{kN\beta} \exp\{\frac{N\beta}{8} \sum_{i=1}^k (E + \frac{x_i}{N})^2\} \int \mathrm{e}^{\frac{k\beta}{8} \sum_{j=1}^{N-k} \lambda_j^2} P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1, \ldots, \mathrm{d}\lambda_{N-k}) \\ &\leq \frac{Z_{N-k}(\alpha - k\beta/4, \beta)}{Z_{N-k}(\alpha, \beta)} C^k \mathrm{e}^{kCb^2} \\ &\leq C^k \mathrm{e}^{kCb^2} \end{aligned}$$

where we used (33) and the fact that for any $x \in [0, 1/2]$, $(1-x)^{-1} \leq 4^x$. □

Hence by (29), (31), (34) and the previous lemma, we have proved that b) of Proposition 6.1 is satisfied. It remains to prove a) for θ given by (7).

4.4. **Preliminary estimates.**

Lemma 4.2 (Bulk eigenvalues)**.** *There is a constant C depending only on the upper bounds on the sequences α and $N\beta$ such that for any $a \in \mathbb{R}$ and $\varepsilon \in (0, 1]$,*

$$P_{\alpha,\beta}^{(N)}(|\lambda_1 - a| \leq \varepsilon) \ \leq \ C\varepsilon \mathrm{e}^{\frac{N\alpha\beta}{2(4\alpha - \beta)} a^2},$$

Proof. We have

$$P_{\alpha,\beta}^{(N)}(|\lambda_1 - a| \leq \varepsilon) \ = \ \frac{1}{Z_N(\alpha, \beta)} \int_{\lambda_1 \in [a \pm \varepsilon]} \mathrm{d}\lambda_1 \mathrm{e}^{-\frac{\alpha}{2}\lambda_1^2} \int_{\mu \in \mathbb{R}^{N-1}} \Delta(\mu)^\beta \prod_j |\lambda_1 - \mu_j|^\beta \mathrm{e}^{-\frac{\alpha}{2} \sum_j \mu_j^2} \mathrm{d}\mu$$

Note that if $|\lambda_1 - a| \leq \varepsilon$, the for any j, $|\lambda_1 - \mu_j| \leq \varepsilon + |\mu_j - a|$. Moreover, for all $x \in \mathbb{R}$, $t > 0$, we have $|x| \leq t^{-1/2} \mathrm{e}^{\frac{tx^2 - 1}{2}}$, hence

$$|x| \leq 2\mathrm{e}^{\frac{x^2}{16}}. \tag{36}$$

Using also the fact that

$$(x + y)^2 \leq 2x^2 + 2y^2, \tag{37}$$

we get that if $|\lambda_1 - a| \leq \varepsilon$,

$$|\lambda_1 - \mu_j|^\beta \leq (\varepsilon + |\mu_j - a|)^\beta \leq 2^\beta \mathrm{e}^{\beta \frac{\varepsilon^2 + (\mu_j - a)^2}{8}}.$$

Hence

$$P_{\alpha,\beta}^{(N)}(|\lambda_1 - a| \leq \varepsilon) \ \leq \ \frac{2^{N\beta} \mathrm{e}^{\frac{N\beta\varepsilon^2}{8}}}{Z_N(\alpha, \beta)} \int_{\lambda_1 \in [a \pm \varepsilon]} \mathrm{d}\lambda_1 \mathrm{e}^{-\frac{\alpha}{2}\lambda_1^2} \int_{\mu \in \mathbb{R}^{N-1}} \Delta(\mu)^\beta \mathrm{e}^{-\frac{\alpha}{2} \sum_j (\mu_j^2 - \beta \frac{(\mu_j - a)^2}{4\alpha})} \mathrm{d}\mu$$

We have

$$\mu_j^2 - \beta\frac{(\mu_j-a)^2}{4\alpha} \;=\; \frac{4\alpha-\beta}{4\alpha}(\mu_j+\frac{\beta a}{4\alpha-\beta})^2 - \frac{\beta}{4\alpha-\beta}a^2$$

Hence

$$\begin{aligned} P^{(N)}_{\alpha,\beta}(|\lambda_1-a|\le\varepsilon) &\le \frac{2^{N\beta}\mathrm{e}^{\frac{N\beta\varepsilon^2}{8}}\mathrm{e}^{\frac{N\alpha\beta}{2(4\alpha-\beta)}a^2}}{Z_N(\alpha,\beta)}\int_{\lambda_1\in[a\pm\varepsilon]}\mathrm{d}\lambda_1\mathrm{e}^{-\frac{\alpha}{2}\lambda_1^2}\int_{\mu\in\mathbb{R}^{N-1}}\Delta(\mu)^\beta\mathrm{e}^{-\frac{\alpha}{2}\sum_j\frac{4\alpha-\beta}{4\alpha}(\mu_j+\frac{\beta a}{4\alpha-\beta})^2}\mathrm{d}\mu \\ &= \varepsilon 2^{N\beta+1}\mathrm{e}^{\frac{N\beta\varepsilon^2}{8}}\mathrm{e}^{\frac{N\alpha\beta}{2(4\alpha-\beta)}a^2}\frac{Z_{N-1}(\alpha-\beta/4,\beta)}{Z_N(\alpha,\beta)} \\ &\le C\varepsilon\mathrm{e}^{\frac{N\alpha\beta}{2(4\alpha-\beta)}a^2}, \end{aligned}$$

we we used (33) to upper bound partition functions quotient. ☐

Lemma 4.3 (Largest eigenvalues)**.** *There is a constant C depending only on the upper bounds on the sequences α and $N\beta$ such that for all $x>0$,*

$$P^{(N)}_{\alpha,\beta}(|\lambda_1|\ge x) \;\le\; C\frac{\mathrm{e}^{-(4\alpha-\beta)\frac{x^2}{8}}}{x}. \tag{38}$$

Proof. We have

$$P^{(N)}_{\alpha,\beta}(|\lambda_1|\ge x) \;=\; \frac{1}{Z_N(\alpha,\beta)}\int_{|\lambda_1|\ge x}\mathrm{d}\lambda_1\mathrm{e}^{-\frac{\alpha}{2}\lambda_1^2}\int_{\mu\in\mathbb{R}^{N-1}}\Delta(\mu)^\beta\prod_j|\lambda_1-\mu_j|^\beta\mathrm{e}^{-\frac{\alpha}{2}\sum_j\mu_j^2}\mathrm{d}\mu$$

Note that by (36) and (37),

$$|\lambda_1-\mu_j|^\beta\le(|\lambda_1|+|\mu_j|)^\beta\le 2^\beta\mathrm{e}^{\beta\frac{\lambda_1^2+\mu_j^2}{8}}.$$

Hence

$$\begin{aligned} P^{(N)}_{\alpha,\beta}(|\lambda_1|\ge x) &\le \frac{2^{N\beta}}{Z_N(\alpha,\beta)}\int_{|\lambda_1|\ge x}\mathrm{d}\lambda_1\mathrm{e}^{-\frac{1}{2}(\alpha-\frac{\beta}{4})\lambda_1^2}\int_{\mu\in\mathbb{R}^{N-1}}\Delta(\mu)^\beta\mathrm{e}^{-\frac{1}{2}(\alpha-\frac{\beta}{4})\sum_j\mu_j^2}\mathrm{d}\mu \\ &= \frac{2^{N\beta}}{\sqrt{\alpha-\beta/4}}\frac{Z_{N-1}(\alpha-\beta/4,\beta)}{Z_N(\alpha,\beta)}\int_{|\lambda_1|\ge x\sqrt{\alpha-\beta/4}}\mathrm{e}^{-\frac{\lambda_1^2}{2}}\mathrm{d}\lambda_1 \end{aligned}$$

Then we conclude using (33) and the fact that for all $y>0$,

$$\int_y^{+\infty}\mathrm{e}^{-\frac{t^2}{2}}\mathrm{d}t \;\le\; \frac{\mathrm{e}^{-\frac{y^2}{2}}}{y}. \tag{39}$$

☐

Lemma 4.4 (Tail of the empirical spectral law)**.** *There are some constants C,c depending only on the upper bounds on the sequences $N\beta,\alpha$ such that for all N and all $x>0$,*

$$P^{(N)}_{\alpha,\beta}(\frac{\lambda_1^2+\cdots\cdots+\lambda_N^2}{N}\ge x) \;\le\; C\mathrm{e}^{-cx}.$$

Proof. We use again the tridiagonal matrix model of (5) for $P^{(N)}_{\alpha,\beta}$ of the λ_i's. We know that $\lambda_1^2+\cdots+\lambda_N^2$ has the same law as $\operatorname{Tr} H^2$, with H the matrix introduced as (5). Note that by the well known convolution relations between Gamma-distributed variables, $\frac{\alpha}{2}\operatorname{Tr} H^2$ has a $\Gamma(\varphi)$-distribution for $\varphi := N(1+(N-1)\beta)/2$. Hence

$$P^{(N)}_{\alpha,\beta}\Big(\frac{\lambda_1^2+\cdots\cdots+\lambda_N^2}{N}\geq x\Big)\ \leq\ \mathbb{P}\Big(G\geq\frac{\alpha N x}{2}\Big),$$

Then, one concludes using the concentration inequalities for Gamma variables (see [9] p. 28-29) which say that for all $u\geq 0$,

$$P(G\geq\varphi(1+u))\ \leq\ \mathrm{e}^{-\varphi(1+u-\sqrt{1+2u})}.$$

□

Lemma 4.5. *For C, c as in the previous lemma, for any u, M, θ such that $cM^2>2\theta\geq 0$,*

$$1\ \leq\ \int \mathrm{e}^{\frac{\theta}{N}\sum_i \log(\frac{|\lambda_i-u|}{M}\vee 1)}P^{(N)}_{\alpha,\beta}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_N)\ \leq\ 1+\frac{2C\theta\mathrm{e}^{cu^2}}{cM^2-2\theta}.$$

Proof. The integral above rewrites

$$1+\theta\int_0^{+\infty}\mathrm{e}^{\theta x}P^{(N)}_{\alpha,\beta}\Big(\frac{1}{N}\sum_i\log\Big(\frac{|\lambda_i-u|}{M}\vee 1\Big)\geq x\Big)\mathrm{d}x. \tag{40}$$

Now, note that as $\log(|\lambda-u|\vee 1)\leq(\lambda-u)^2$,

$$\begin{aligned}P^{(N)}_{\alpha,\beta}\Big(\frac{1}{N}\sum_i\log\Big(\frac{|\lambda_i-u|}{M}\vee 1\Big)\ \geq\ x\Big)\ &\leq\ P^{(N)}_{\alpha,\beta}\Big(\frac{(\lambda_1-u)^2+\cdots\cdots+(\lambda_N-u)^2}{N}\geq M^2x\Big)\\ &\leq\ P^{(N)}_{\alpha,\beta}\Big(\frac{2\lambda_1^2+2u^2+\cdots\cdots+2\lambda_N^2+2u^2}{N}\geq M^2x\Big)\\ &\leq\ P^{(N)}_{\alpha,\beta}\Big(\frac{\lambda_1^2+\cdots\cdots+\lambda_N^2}{N}\geq\frac{M^2x}{2}-u^2\Big).\end{aligned}$$

Then one concludes using (40) and the previous lemma. □

4.5. **Convergence of the correlation functions.**

Let us now prove a) of Proposition 6.1 for θ given by (7). Note first that by b) of Proposition 2.1, we know that

$$\int\log|E-x|\mathrm{d}\mu_\gamma(x)\ <\ \infty. \tag{41}$$

Besides, by (27), (30), (32) and (35), it suffices to prove that for each k and each $x_1,\ldots,x_k\in\mathbb{R}$, the quantity $\widetilde{R}^{(N)}_k(x_1,\ldots,x_k)$ defined at (28) satisfies, as $N\to\infty$,

$$\widetilde{R}^{(N)}_k(x_1,\ldots,x_k)\ \underset{N\to\infty}{\longrightarrow}\ \exp\Big\{2\gamma k\int\log|E-x|\mathrm{d}\mu_\gamma(x)\Big\}\,. \tag{42}$$

4.5.1. *Upper-bound.* Let us prove that for any fixed k and $x_1, \dots, x_k$,

$$\limsup_{N\to\infty} \widetilde{R}_k^{(N)}(x_1,\dots,x_k) \;\le\; \exp\left\{2\gamma k \int \log|E-x|\mathrm{d}\mu_\gamma(x)\right\}. \tag{43}$$

For $\varepsilon > 0$, set

$$\widetilde{R}_k^{(N,\varepsilon)}(x_1,\dots,x_k) \;:=\; \int \mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log(|E+\frac{x_i}{N}-\lambda_j|\vee\varepsilon)} P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\dots,\mathrm{d}\lambda_{N-k}).$$

As $\widetilde{R}_k^{(N)} \le \widetilde{R}_k^{(N,\varepsilon)}$ and, by (41), $\inf_{\varepsilon>0}\int \log(|x-E|\vee\varepsilon)\mathrm{d}\mu_\gamma(x) = \int \log|x-E|\mathrm{d}\mu_\gamma(x)$, it suffices to prove that for any $\varepsilon > 0$ small enough, we have

$$\limsup_{N\to\infty} \widetilde{R}_k^{(N,\varepsilon)}(x_1,\dots,x_k) \;\le\; \exp\left\{2\gamma k \int \log(|E-x|\vee\varepsilon)\mathrm{d}\mu_\gamma(x)\right\}. \tag{44}$$

Note now that for any $M > 0$ large enough, as

$$\log(|E+x_i/N-x|\vee\varepsilon) = \log\{(|E+x_i/N-x|\vee\varepsilon)\wedge M\} + \log\left(\frac{|E+x_i/N-x|}{M}\vee 1\right)$$

the function $\widetilde{R}_k^{(N,\varepsilon)}(x_1,\dots,x_k)$ rewrites

$$\begin{aligned}
&\widetilde{R}_k^{(N,\varepsilon)}(x_1,\dots,x_k)\\
&= \int \mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log\{(|E+\frac{x_i}{N}-\lambda_j|\vee\varepsilon)\wedge M\}}\mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log\left(\frac{|E+\frac{x_i}{N}-\lambda_j|}{M}\vee 1\right)} P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\dots,\mathrm{d}\lambda_{N-k})\\
&= \int \mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log\{(|E+\frac{x_i}{N}-\lambda_j|\vee\varepsilon)\wedge M\}} P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\dots,\mathrm{d}\lambda_{N-k}) +\\
&\quad \int \underbrace{\mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log\{(|E+\frac{x_i}{N}-\lambda_j|\vee\varepsilon)\wedge M\}}}_{:=X}\underbrace{\left(\mathrm{e}^{\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log\left(\frac{|E+\frac{x_i}{N}-\lambda_j|}{M}\vee 1\right)}-1\right)}_{:=Y} P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\dots,\mathrm{d}\lambda_{N-k})
\end{aligned}$$

By Proposition 2.1, we know that under the law $P_{\alpha,\beta}^{(N-k)}$, the L^∞-bounded sequence of random variables $\beta\sum_{i=1}^k\sum_{j=1}^{N-k}\log\{(|E+\frac{x_i}{N}-\lambda_j|\vee\varepsilon)\wedge M\}$ converges in probability, as $N\to\infty$, to

$$2\gamma k\int \log\{(|E-x|\vee\varepsilon)\wedge M\}\mathrm{d}\mu_\gamma(x)$$

(one gets rid of the $\frac{x_i}{N}$'s by noticing, for example, that the convolution of probability measures is continuous with respect to the weak topology and that $\delta_{\lambda_i/N}$ converges to δ_0).

Note that by choosing M large enough, one can make $\int \log\{(|E-x|\vee\varepsilon)\wedge M\}\mathrm{d}\mu_\gamma(x)$ as close as we want from $\int\log(|E-x|\vee\varepsilon)\mathrm{d}\mu_\gamma(x)$. Moreover, one can easily adapt the proof of Lemma 4.1 to see that $\int X^2 P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_{N-k})$ is bounded by a constant independent of M, hence by Cauchy-Schwartz, to prove (44), it suffices to prove that $\int Y^2 P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_{N-k})$ can be made as small as we want if M is large enough. Note that for

$$Y_i := \mathrm{e}^{\beta\sum_{j=1}^{N-k}\log\left(\frac{|E+\frac{x_i}{N}-\lambda_j|}{M}\vee 1\right)},$$

we have $Y = Y_1\cdots Y_k - 1$. Thus by the Hölder inequality, it is enough to prove that for M large enough, earh Y_i can have its k-th and $2k$-th moment as close as we want from 1, which is a direct consequence of Lemma 4.5.

4.5.2. *Lower bound.* To obtain the analogous lower bound

$$\liminf_{N\to\infty}\widetilde{R}_k^{(N)}(x_1,\ldots,x_k)\ \geq\ \exp\left\{2\gamma k\int\log|E-x|\mathrm{d}\mu_\gamma(x)\right\}, \tag{45}$$

we observe that first by Jensen's inequality and then by exchangeability,

$$\begin{aligned}\log\widetilde{R}_k^{(N)}(x_1,\ldots,x_k)\ &\geq\ \int\beta\sum_{i=1}^{k}\sum_{j=1}^{N-k}\log|E+\frac{x_i}{N}-\lambda_j|P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_{N-k})\\ &=\ (N-k)\beta\int\sum_{i=1}^{k}\log|E+\frac{x_i}{N}-\lambda_1|P_{\alpha,\beta}^{(N-k)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_{N-k})\end{aligned}$$

Hence as $(N-k)\beta\longrightarrow 2\gamma$ and the triplet $(N,\beta(N),\alpha(N))$ satisfies the same hypotheses as $(N-k,\beta(N),\alpha(N))$, it suffices to prove that for any fixed x, we have

$$\liminf_{N\to\infty}\int\log|E+\frac{x}{N}-\lambda_1|P_{\alpha,\beta}^{(N)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_N)\ \geq\ \int\log|E-t|\mathrm{d}\mu_\gamma(t).$$

As, by exchangeability, μ_γ is also the weak limit of the distribution of λ_1 under $P_{\alpha,\beta}^{(N)}$, we know that for any $\varepsilon>0$,

$$\liminf_{N\to\infty}\int\log(|E+\frac{x}{N}-\lambda_1|\vee\varepsilon)P_{\alpha,\beta}^{(N)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_N)\ \geq\ \int\log|E-t|\mathrm{d}\mu_\gamma(t)$$

(and one can get rid of $\frac{x}{N}$ for the same reason as in Section 4.5.1 above). Hence it suffices that for ε small enough,

$$\limsup_{N\to\infty}\left|\int\left(\log(|E+\frac{x}{N}-\lambda_1|\vee\varepsilon)-\log|E+\frac{x}{N}-\lambda_1|\right)P_{\alpha,\beta}^{(N)}(\mathrm{d}\lambda_1,\ldots,\mathrm{d}\lambda_N)\right|$$

can be made as small as desired. But for any random variable $X>0$,

$$\mathbb{E}[\log(X\vee\varepsilon)-\log(X)]=\mathbb{E}[(\log\varepsilon-\log(X))\mathbb{1}_{X\leq\varepsilon}]=\int_0^\varepsilon\frac{\mathbb{P}(X\leq t)}{t}\mathrm{d}t.$$

Here, by Lemma 4.2, there is a constant C such that

$$P_{\alpha,\beta}^{(N)}(|E+\frac{x}{N}-\lambda_1|\leq t)\ \leq\ Ct,$$

which allows to get the desired bound.

5. CASE WHERE $N^{-1} \ll \beta \ll 1/\log(N)$

5.1. **Partition functions: proofs of (18) and (19).** It follows from (33) that for all N, for all $1 \le k \le N$,

$$\begin{aligned}\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} &= \alpha^{\frac{k}{2}}\alpha^{\frac{\beta}{4}(N(N-1)-(N-k)(N-k-1))}(2\pi)^{-k/2}\Gamma(1+\frac{\beta}{2})^k\prod_{\ell=0}^{k-1}\frac{1}{\Gamma(1+(N-\ell)\beta/2)}\\ &= \alpha^{\frac{k}{2}}\alpha^{\frac{\beta}{4}(2kN-k^2+k)}(2\pi)^{-k/2}\Gamma(1+\frac{\beta}{2})^k\prod_{\ell=0}^{k-1}\frac{1}{\Gamma(1+(N-\ell)\beta/2)}\end{aligned}$$

But by the Stirling formula, for $z \to +\infty$,

$$\Gamma(1+z) \sim \sqrt{2\pi z}\exp\left\{z\log(z)-z\right\}. \tag{46}$$

Note that our hypothesis on β implies that for any fixed $\ell = 0,\dots,k-1$, we have

$$(N-\ell)\beta/2\log((N-\ell)\beta/2)-(N-\ell)\beta/2 = N\beta/2\log(N\beta/2)-N\beta/2+o(1),$$

so that, as (11) implies that $N\beta\log\frac{2\alpha}{N\beta} \ll 1$ and $\beta\log\alpha \ll 1$,

$$\frac{Z_{N-k}(\alpha,\beta)}{Z_N(\alpha,\beta)} \sim \left(\frac{\mathrm{e}^{N\beta/2}}{2\pi}\right)^k \times \left(\exp\left\{\frac{N\beta}{2}\left[\log\alpha-\log(N\beta/2)\right]\right\}\right)^k$$

By (11), we get (18). The upper bound (19) comes in the same way, noticing that the error in (46) is uniformly bounded on $z \ge 0$.

5.2. **Tail estimate.**

Lemma 5.1. *Let $\beta = \beta(N)$ and $\alpha = \alpha(N)$ be satisfying (11). Then there is a constant C depending only on the sequences α and β such that for all N, for all $x > 0$,*

$$P^{(N)}_{\alpha,\beta}(|\lambda_1| \ge x) \le C\mathrm{e}^{\left(-\frac{x^2}{2}+C\right)N\beta/2}. \tag{47}$$

Proof. We have

$$P^{(N)}_{\alpha,\beta}(|\lambda_1| \ge x) = \frac{1}{Z_N(\alpha,\beta)}\int_{|\lambda_1|\ge x}\mathrm{d}\lambda_1\mathrm{e}^{-\frac{\alpha}{2}\lambda_1^2}\int_{\mu\in\mathbb{R}^{N-1}}\Delta(\mu)^\beta\prod_j|\lambda_1-\mu_j|^\beta\mathrm{e}^{-\frac{\alpha}{2}\sum_j\mu_j^2}\mathrm{d}\mu$$

Note that by (36) and (37),

$$|\lambda_1-\mu_j|^\beta \le (|\lambda_1|+|\mu_j|)^\beta \le 2^\beta\mathrm{e}^{\beta\frac{\lambda_1^2+\mu_j^2}{8}}.$$

Hence

$$\begin{aligned}
P^{(N)}_{\alpha,\beta}(|\lambda_1|\ge x) &\le \frac{2^{N\beta}}{Z_N(\alpha,\beta)}\int_{|\lambda_1|\ge x}\mathrm{d}\lambda_1 \mathrm{e}^{-\frac{\alpha-\beta/4}{2}\lambda_1^2}\int_{\mu\in\mathbb{R}^{N-1}}\Delta(\mu)^\beta \mathrm{e}^{-\frac{\alpha-\beta/4}{2}\sum_j\mu_j^2}\mathrm{d}\mu\\
&= \frac{2^{N\beta}}{Z_N(\alpha,\beta)\sqrt{\alpha-\beta/4}}\int_{|\lambda_1|\ge x\sqrt{\alpha-\beta/4}}\mathrm{e}^{-\frac{\lambda_1^2}{2}}\mathrm{d}\lambda_1\times Z_{N-1}(\alpha-\frac{\beta}{4},\beta)\\
&\le \frac{2^{N\beta+1}\mathrm{e}^{-\frac{x^2(\alpha-\beta/4)}{2}}}{x(\alpha-\beta/4)}\frac{Z_{N-1}(\alpha-\frac{\beta}{4},\beta)}{Z_N(\alpha,\beta)}\\
&= \frac{2^{N\beta+1}\mathrm{e}^{-\frac{x^2(\alpha-\beta/4)}{2}}}{x(\alpha-\beta/4)}(1-\beta/(4\alpha))^{-(\frac{(N-1)(N-2)}{4}\beta+\frac{N-1}{2})}\frac{Z_{N-1}(\alpha,\beta)}{Z_N(\alpha,\beta)}
\end{aligned}$$

Let us now use (19) and for example the fact that $(1-y)^{-1}\le \mathrm{e}^{2y}$ when $y\in[0,1/2]$. We get

$$P^{(N)}_{\alpha,\beta}(|\lambda_1|\ge x) \le \frac{M}{2\pi}\frac{1}{x\alpha}\exp\{-\frac{x^2}{2}(\alpha-\beta/4)+N^2\beta^2/(8\alpha)+N\beta/(2\alpha)+N\beta/2\},$$

which allows to conclude, as we already noticed that (11) implies that $2\alpha\sim N\beta\gg 1$. $\square$

5.3. **Proof of Theorem 2.5.** We first define the probability measure on $\mathbb{R}^N$

$$P^{(N,h)}_{\alpha,\beta}(\mathrm{d}x) := \underbrace{\frac{1}{Z^h_N(\alpha,\beta)}\Delta(x)^\beta\exp\{-\frac{\alpha}{2}\sum_{i=1}^N x_i^2+\beta\sum_{i=1}^N h(x_i)\}}_{:=\rho^{(N,h)}_{\alpha,\beta}(x_1,\ldots,x_N)}\mathrm{d}x_1\cdots\mathrm{d}x_N, \tag{48}$$

where

$$Z^h_N(\alpha,\beta) = \int\Delta(x)^\beta \mathrm{e}^{-\frac{\alpha}{2}\sum_{i=1}^N x_i^2+\beta\sum_{i=1}^N h(x_i)}\mathrm{d}x_1\cdots\mathrm{d}x_N \tag{49}$$

is the normalisation constant. Let, for $i=1,2$,

$$u^{i,h}_N(x_1,\ldots,x_i) := \int_{x_{i+1},\ldots,x_N}\rho^{(N,h)}_{\alpha,\beta}(x_1,\ldots,x_N)\mathrm{d}x_{i+1}\cdots\mathrm{d}x_N$$

be the i-th correlation function of $\rho^{(N,h)}_{\alpha,\beta}$.

Lemma 5.2. *Let $\psi:\mathbb{R}\to\mathbb{C}$ be a $\mathcal{C}^1$ function on $\mathbb{R}$ such that the real and imaginary parts of ψ' are bounded below. Then we have*

$$\begin{aligned}
\beta\frac{N-1}{2}\iint\frac{\psi(t)-\psi(s)}{t-s}u^{2,h}_N(s,t)\mathrm{d}s\mathrm{d}t+\int_{\mathbb{R}}(\beta h'(t)-\alpha t)\psi(t)u^{1,h}_N(t)\mathrm{d}t&\\
+\int_{\mathbb{R}}\psi'(t)u^{1,h}_N(t)\mathrm{d}t &= 0.
\end{aligned}\tag{50}$$

Proof. As (50) is linear in ψ, one can suppose ψ to be real-valued. Then for $\theta \geq 0$ small enough, the function $y + \theta\psi(y)$ is an homeomorphism on $\mathbb{R}$, hence one can make the change of variable $x_i = y_i + \theta\psi(y_i)$ in (49). We get

$$\begin{aligned} Z_N^h &= \int_{y\in\mathbb{R}^N} \prod_{1\leq i<j\leq N} |y_j - y_i + \theta(\psi(y_j) - \psi(y_i))|^\beta \qquad (51) \\ & \quad \mathrm{e}^{-\frac{\alpha}{2}\sum_{i=1}^N (y_i^2 + 2\theta y_i\psi(y_i) + \theta^2\psi(y_i)^2) + \beta\sum_{i=1}^N h(y_i+\theta\psi(y_i))} \prod_{i=1}^N (1+\theta\psi'(y_i))\mathrm{d}y_1\cdots\mathrm{d}y_N \end{aligned}$$

Let us compute the derivative, with respect to θ, at $\theta = 0$, of the RHT of (51). We have

$$\partial_{\theta,\theta=0}|y_j - y_i + \theta(\psi(y_j) - \psi(y_i))|^\beta = \beta\frac{\psi(y_j) - \psi(y_i)}{y_j - y_i}|y_j - y_i|^\beta,$$

we also have

$$\partial_{\theta,\theta=0}\mathrm{e}^{-\frac{\alpha}{2}\sum_{i=1}^N (y_i^2 + 2\theta y_i\psi(y_i) + \theta^2\psi(y_i)^2)} = -\alpha\sum_{i=1}^N y_i\psi(y_i),$$

$$\partial_{\theta,\theta=0}\mathrm{e}^{\beta\sum_{i=1}^N h(y_i+\theta\psi(y_i))} = \beta\sum_{i=1}^N h'(y_i)\psi(y_i),$$

and

$$\partial_{\theta,\theta=0}\prod_{i=1}^N (1+\theta\psi'(y_i)) = \sum_{i=1}^N \psi'(y_i).$$

Hence

$$\begin{aligned} & \frac{\partial_{\theta,\theta=0}(\text{RHT of (51)})}{Z_N^h} \\ = \; & \beta\sum_{1\leq i<j\leq N}\int_{y\in\mathbb{R}^N} \frac{\psi(y_j) - \psi(y_i)}{y_j - y_i}\rho_N^h(y_1,\ldots,y_N)\mathrm{d}y_1\cdots\mathrm{d}y_N \\ & -\alpha\sum_{i=1}^N\int_{y\in\mathbb{R}^N} y_i\psi(y_i)\rho_N^h(y_1,\ldots,y_N)\mathrm{d}y_1\cdots\mathrm{d}y_N \\ & +\beta\sum_{i=1}^N\int_{y\in\mathbb{R}^N} h'(y_i)\psi(y_i)\rho_N^h(y_1,\ldots,y_N)\mathrm{d}y_1\cdots\mathrm{d}y_N \\ & +\sum_{i=1}^N\int_{y\in\mathbb{R}^N} \psi'(y_i)\rho_N^h(y_1,\ldots,y_N)\mathrm{d}y_1\cdots\mathrm{d}y_N \end{aligned}$$

We get exactly (50). □

Now, we define, for z such that $\Im z > 0$,

$$H_N(z) := \int \frac{h'(t)}{z-t} u_N^{1,h}(t)\mathrm{d}t \qquad ; \qquad U_N(z) := \int \frac{1}{z-t} u_N^{1,h}(t)\mathrm{d}t$$
$$k_N(s,t) := N u_N^{1,h}(s) u_N^{1,h}(t) - (N-1) u_N^{2,h}(s,t)$$

and

$$\begin{aligned} K_N(z) &:= N \iint \left(\frac{1}{(z-t)(z-s)} - \frac{1}{2(z-t)^2} - \frac{1}{2(z-s)^2} \right) k_N(s,t)\mathrm{d}t\mathrm{d}s \qquad (52) \\ &= -\frac{1}{2} \iint \frac{(t-s)^2}{(z-t)^2(z-s)^2} k_N(s,t)\mathrm{d}s\mathrm{d}t. \end{aligned}$$

We also introduce

$$U(z) := \frac{1}{2}(z - \sqrt{z^2-4}) \qquad\qquad (\Im(z) > 0), \qquad (53)$$

where when $\Re(z) \geq 0$ (resp. ≤ 0) , $\sqrt{z^2-4}$ is computed with the determination of the square root on $\mathbb{C}\backslash(-\infty, 0)$ (resp. on $\mathbb{C}\backslash(0, +\infty)$) with positive values on $[0, +\infty)$ (resp. such that $\sqrt{-1} = \mathrm{i}$). It is well known that U is the Stieltjes transform of the semicircle law σ.

Lemma 5.3. *On the upper half-plane, we have*

$$N\beta(U_N - U)(2U - \frac{2\alpha}{N\beta} z + U_N - U) = \Delta_N := \beta N^{-1} K_N - 2\beta H_N + (2-\beta) U_N' + (2\alpha - N\beta) U^2. \qquad (54)$$

Proof. We shall apply the previous lemma with $\psi(t) = \frac{1}{z-t}$. Note that $\frac{\psi(t)-\psi(s)}{t-s} = \frac{1}{(z-t)(z-s)}$, so that we have :

$$\begin{aligned} & N(N-1) \iint \frac{\psi(t)-\psi(s)}{t-s} u_N^{2,h}(s,t)\mathrm{d}s\mathrm{d}t \\ = & \iint \frac{1}{(z-t)(z-s)} N(N-1) u_N^{2,h}(s,t)\mathrm{d}s\mathrm{d}t \\ = & \iint \frac{1}{(z-t)(z-s)} (-N k_N(s,t) + N^2 u_N^{1,h}(s) u_N^{1,h}(t))\mathrm{d}s\mathrm{d}t \\ = & N^2 U_N(z)^2 - N \iint \frac{1}{(z-t)(z-s)} k_N(s,t)\mathrm{d}s\mathrm{d}t \\ = & N^2 U_N(z)^2 - N \iint \left(\frac{1}{(z-t)(z-s)} - \frac{1}{2(z-t)^2} - \frac{1}{2(z-s)^2} \right) k_N(s,t)\mathrm{d}s\mathrm{d}t \\ & -N \iint \frac{1}{(z-t)^2} k_N(s,t)\mathrm{d}s\mathrm{d}t \\ = & N^2 U_N(z)^2 - K_N(z) + n U_N'(z) \end{aligned}$$

(where we use the fact that for any function $f(t)$, $\iint f(t)k_N(s,t)\mathrm{d}s\mathrm{d}t = \int f(t)u_N^{1,h}(t)\mathrm{d}t$). We also have

$$\int -t\psi(t)u_N^{1,h}(t)\mathrm{d}t = 1 - zU_N(z) \quad ; \quad \int \psi'(t)u_N^{1,h}(t)\mathrm{d}t = -U_N'(z),$$

so $2N^{-2}\beta^{-1}\times$(50) rewrites

$$U_N(z)^2 - N^{-2}K_N(z) + N^{-1}U_N'(z) + 2N^{-1}H_N(z) + \frac{2\alpha}{N\beta}(1 - zU_N(z)) - \frac{2}{N\beta}U_N'(z) = 0$$

i.e.

$$U_N(z)^2 - \frac{2\alpha}{N\beta}zU_N(z) + \frac{2\alpha}{N\beta} = N^{-2}K_N(z) - 2N^{-1}H_N(z) + (\frac{2}{N\beta} - \frac{1}{N})U_N'(z). \tag{55}$$

One gets (54), using the well known equation $U(z)^2 - zU(z) + 1 = 0$ (see [6, Eq. (2.4.6)]). $\square$

A key step in the proof of the theorem will be to prove that as $N \to \infty$,

$$\beta K_N(z) \ll N. \tag{56}$$

We shall now prove (56). Let $(y_1, \dots, y_N)$ be a random vector with distribution $P_{\alpha,\beta}^{(N,h)}$ and for $g \in \mathcal{C}_b(\mathbb{R}, \mathbb{C})$, define the random variable

$$\hat{\mu}_N^h(g) := \sum_{i=1}^N g(y_i),$$

with variance $\Sigma_N(g) := \mathbb{E}[|\hat{\mu}_N^h(g) - \mathbb{E}[\hat{\mu}_N^h(g)]|^2]$. As

$$K_N(z) = \mathbb{E}[(\hat{\mu}_N^h(g) - \mathbb{E}[\hat{\mu}_N^h(g)])^2] \tag{57}$$

for $g(t) = (z-t)^{-1}$, we have

$$|K_N(z)| \le \Sigma_N(\frac{1}{z-t}). \tag{58}$$

Note that as

$$\begin{aligned} \mathbb{E}[\hat{\mu}_N^h(g)] &= N\int u_N^{1,h}(t)g(t)\mathrm{d}t \\ \mathbb{E}[|\hat{\mu}_N^h(g)|^2] &= \sum_{i,j=1}^N \mathbb{E}[g(y_i)\overline{g(y_j)}] \\ &= N(N-1)\iint g(s)\overline{g(t)}u_N^{2,h}(s,t)\mathrm{d}s\mathrm{d}t + N\int u_N^{1,h}(t)|g(t)|^2\mathrm{d}t, \end{aligned}$$

we deduce

$$\begin{aligned}\Sigma_N(g) &= N(N-1)\iint g(s)\overline{g(t)}(u_N^{2,h}(s,t)-u_N^{1,h}(s)u_N^{1,h}(t))\mathrm{d}s\mathrm{d}t\\ &\quad +N\left(\int u_N^{1,h}(t)|g(t)|^2\mathrm{d}t-|\int u_N^{1,h}(t)g(t)\mathrm{d}t|^2\right)\end{aligned}$$

Lemma 5.4. *There is $L>0$ and $c>0$ such that for any fixed function g,*

$$\begin{aligned}\Sigma_N(g) &\le N(N-1)\iint_{[-L,L]^2} g(s)\overline{g(t)}(u_N^{2,h}(s,t)-u_N^{1,h}(s)u_N^{1,h}(t))\mathrm{d}s\mathrm{d}t\\ &\quad +N\left(\int_{-L}^{L} u_N^{1,h}(t)|g(t)|^2\mathrm{d}t-|\int_{-L}^{L} u_N^{1,h}(t)g(t)\mathrm{d}t|^2\right)+4N^2\|g\|_\infty \mathrm{e}^{-cN\beta}\end{aligned}$$

Proof. Using the fact that $|\sum_{i=1}^N h(x_i)|\le N\|h\|_\infty$, we see that the probability measure $P_{\alpha,\beta}^{(N,h)}$ defined at (48) and its normalization constant can be controlled thanks to the probability measure $P_{\alpha,\beta}^{(N)}$ and its normalization constant: for any Borel set $A\subset\mathbb{R}^N$, we have

$$P_{\alpha,\beta}^{(N,h)}(A)\le \mathrm{e}^{2N\beta\|h\|_\infty}P_{\alpha,\beta}^{(N)}(A).$$

It follows that up to a change of the constant C, Lemma 5.1 is also true for $P_{\alpha,\beta}^{(N,h)}$, which allows to conclude. ☐

This lemma allows to reduce the problem to a compact set, and after rescaling, one can turn the compact set in question to $[-1/2,1/2]$: we deduce, as in [17], that for $w:=z/L$ and

$$\rho_N^2(t,s):=u_N^{2,h}(Lt,Ls),\qquad \rho_N^1(t):=u_N^{1,h}(Lt).$$

we have

$$\Sigma_N(\frac{1}{z-t})\le N(N-1)\int_{-1/2}^{1/2}\frac{1}{w-t}G_N(x\mapsto\frac{1}{\overline{w}-x})(t)\mathrm{d}t+4N^2e^{-cN\beta}(\Im z)^{-2}, \tag{59}$$

where G_N is the operator on $L^2([-1/2,1/2],\mathrm{d}x)$ defined by

$$G_N(f)(t)=\int f(s)(\rho_N^2(t,s)-\rho_N^1(t)\rho_N^1(s))\mathrm{d}s+\frac{\rho_N(t)}{N-1}\left(\frac{1}{L}f(t)-\int_{-1/2}^{1/2}\rho_N(s)f(s)\mathrm{d}s\right).$$

Thus to prove the estimate of interest (56), we have to upper bound:

$$N(N-1)\int_{-1/2}^{1/2}\frac{1}{w-t}G_N(x\mapsto\frac{1}{\overline{w}-x})(t)\mathrm{d}t. \tag{60}$$

Following [17], we introduce the integral operator P_w on $L^2([-1/2,1/2],\mathrm{d}x)$ with kernel $P_w(t,s)=\frac{1}{(w-t)(\overline{w}-s)}$. Then $(s,t)\longmapsto\frac{1}{w-t}G_N(x\mapsto\frac{1}{\overline{w}-x})(s)$ is an integral operator satisfying the hypothesis of Theorem 2.12 of [24]. This trace class operator is nothing but $P_w G_N$, thus by this theorem, we have

$$\int_{-1/2}^{1/2} \frac{1}{w-t} G_N(\frac{1}{\overline{w}-s})(t)\mathrm{d}t = \mathrm{Tr}(P_w G_N). \tag{61}$$

We will not here recall all the arguments used in [17] Lemma 3.12 and Proposition 3.9 to estimate this trace. The proof transfers to our setting using minor modifications (essentially replacing N by $N\beta$). Note that the important feature of h is that is Lipschitz on compact sets in our case. Thus we simply state the final estimate we will use in this article, namely the following lemma :

Lemma 5.5. *We have* $\mathrm{Tr}(P_w G_N) \le C N^{-1}\log(N)$ *for some constant* C.

It follows, by (58), (59) and (61), that

$$|\beta K_N(z)| \le \beta \Sigma_N(\frac{1}{z-t}) \le CN\beta\log(N) + +4N^2\beta \mathrm{e}^{-cN\beta}(\Im z)^{-2}.$$

This is of course $\ll N$, so the estimate of interest (56) is proved.

As $\beta\log(N) \ll 1$ and $N\beta - 2\alpha \ll 1$, by (54), we deduce that, uniformly on compact subsets of $\mathbb{C}^+$, one has that

$$N\beta(U_N(z) - U(z)) \longrightarrow \frac{2U'(z)}{2U(z)-z} = \frac{z}{z^2-4} - \frac{1}{\sqrt{z^2-4}}. \tag{62}$$

One recognizes easily that the RHT of (62) is the Stieltjes transform of the null mass signed measure ν of (13) (to do that, use the fact that $U(z)$, given by (53), is the Stieltjes transform of the semi-circle law and then use an integration by parts).

The rest of the proof of the theorem is an easy adaptation of the proof of Theorem 2.4 in [17] (p. 169-172). The main idea is to define

$$F(\lambda) := \log \mathbb{E}\, \mathrm{e}^{\beta\lambda\sum_{j=1}^N h(x_j)} - N\beta\lambda \int h(t)\mathrm{d}\sigma(t),$$

to notice that

$$\partial_\lambda F(\lambda) = N\int_{\mathbb{R}} h(t) u_N^{1,\lambda h}(t)\mathrm{d}t,$$

to prove (12) for derivatives and to deduce (12) by dominated convergence. We use (62), namely the convergence

$$N\beta \int \frac{1}{z-t}\left(u_N^{1,h}(t) - \mathbb{1}_{|t|\le 2}\frac{\sqrt{4-t^2}}{2\pi}\right)\mathrm{d}t \longrightarrow \int \frac{1}{z-t}\mathrm{d}\nu(t),$$

in Fourier transform manipulations, precisely via the formula

$$\int_0^\infty \hat{\delta}_N(\xi)\mathrm{e}^{\mathrm{i}\xi z}\mathrm{d}\xi = \mathrm{i}\int \frac{1}{z-t}\delta_N(t)\mathrm{d}t \qquad (\Im z > 0)$$

with $\delta_N(t) := N\beta(u_N^{1,h}(t) - \mathbb{1}_{|t|\le 2}\frac{\sqrt{4-t^2}}{2\pi})$.

6. Appendix: Poisson limit for point processes

Let $\mathcal{X}$ be a locally compact Polish space and μ be a Radon measure on $\mathcal{X}$. We consider an exchangeable random vector $(\lambda_1, \ldots, \lambda_N)$ taking values on $\mathcal{X}$ implicitly depending on N, with density $\rho^{(N)}$ with respect to $\mu^{\otimes N}$. We define, for $1 \le k \le N$, the k-th correlation function on $\mathcal{X}^k$ by the formula

$$R_k^{(N)}(x_1, \ldots, x_k) := \frac{N!}{(N-k)!} \int_{(x_{k+1},\ldots,x_N)\in\mathcal{X}^{N-k}} \rho^{(N)}(x_1, \ldots, x_N)\mathrm{d}\mu^{\otimes N-k}(x_{k+1}, \ldots, x_N).$$

Proposition 6.1. *Suppose that there is $\theta \ge 0$ independent of N such that the correlation functions $R_k^{(N)}$ satisfy:*

a) *For each $k \ge 1$, on $\mathcal{X}^k$, we have the pointwise convergence*

$$R_k^{(N)}(x_1, \ldots, x_k) \underset{N\to\infty}{\longrightarrow} \theta^k, \tag{63}$$

b) *For each compact $\mathcal{K} \subset \mathcal{X}$, there is $\Theta_{\mathcal{K}}$ such that for all k, N, on $\mathcal{K}^k$, we have*

$$\mathbb{1}_{k\le N} R_k^{(N)}(x_1, \ldots, x_k) \;\le\; \Theta_{\mathcal{K}}^k \tag{64}$$

Then the point process $\sum_{i=1}^N \delta_{\lambda_i}$ converges in distribution to a Poisson point process with intensity $\theta\mathrm{d}\mu$ as $N \to \infty$.

Proof. Note that the Poisson point process M with intensity $\theta\mathrm{d}\mu$ is characterized, among random random Radon measures on $\mathcal{X}$, by the fact that for any compactly supported continuous function f on $\mathcal{X}$, we have

$$\mathbb{E}\,\mathrm{e}^{\langle M,f\rangle} = \exp\left(\theta \int (\mathrm{e}^{f(x)} - 1)\mathrm{d}\mu(x)\right).$$

So let us fix f a compactly supported continuous function on $\mathcal{X}$. Then, with the convention $R_0^{(N)} = 1$,

$$\begin{aligned}
\mathbb{E}\,\mathrm{e}^{\sum_{i=1}^N f(\lambda_i)} &= \mathbb{E}\prod_{i=1}^{N}(1+(\mathrm{e}^{f(\lambda_i)}-1)) \\
&= \sum_{P\subset\{1,\dots,n\}} \mathbb{E}\prod_{i\in P}(\mathrm{e}^{f(\lambda_i)}-1) \\
&= \sum_{k=0}^{N}\binom{N}{k}\mathbb{E}\prod_{i=1}^{k}(\mathrm{e}^{f(\lambda_i)}-1) \\
&= \sum_{k=0}^{N}\frac{1}{k!}\int\prod_{i=1}^{k}(\mathrm{e}^{f(x_i)}-1)R_k^{(N)}(x_1,\dots,x_k)\mathrm{d}\mu^{\otimes k}(x_1,\dots,x_k)
\end{aligned}$$

This proves the proposition. □

References

[1] R. Allez, J.-P. Bouchaud, A. Guionnet *Invariant Beta ensembles and the Gauss-Wigner crossover* Physical review letters 109 (9), 094–102.

[2] R. Allez, J.-P. Bouchaud, S.N. Majumdar, P. Vivo *Invariant β-Wishart ensembles, crossover densities and asymptotic corrections to the Marcenko-Pastur law* Journal of Physics A: Mathematical and Theoretical 46 (1), 015001

[3] R. Allez, A Guionnet *A diffusive model for invariant β-ensembles* Electron. J. Probab 18 (62), 1–30.

[4] R. Allez, L. Dumaz *From sine kernel to Poisson statistics* Elec. J. Probab. (2014), Vol. 19, 1–25.

[5] R. Allez, L. Dumaz *Tracy–Widom at High Temperature* J. Stat. Phys. (2014), Vol. 156, No 6, 1146–1183.

[6] G. Anderson, A. Guionnet, O. Zeitouni *An Introduction to Random Matrices.* Cambridge studies in advanced mathematics, 118 (2009).

[7] Z. Bao, Z. Su *Local Semicircle law and Gaussian fluctuation for Hermite β-ensemble*, arXiv.

[8] G. Ben Arous, A. Guionnet. *Large deviations for Wigner's law and Voiculescu's non commutative entropy*, Probab. Theory Related Fields, **108**, (1997) 517–542.

[9] S. Boucheron, G. Lugosi, P. Massart *Concentration inequalities*, Oxford, 2013.

[10] P. Bourgade *Bulk universality for one-dimensional log-gases*, proceedings of the XVIIth International Congress On Mathematical Physics, World Scientific Publishing, 2013.

[11] P. Bourgade, L. Erdös, H.T. Yau *Bulk universality of general β-ensembles with non-convex potential*, J. of Math. Phys., special issue in honor of E. Lieb's 80th birthday, Vol. 53, 2012.

[12] P. Bourgade, L. Erdös, H.T. Yau *Universality of general β-ensembles*, Duke Math. J., Vol. 163, no. 6, 1127–1190, 2014.

[13] P. Bourgade, L. Erdös, H.T. Yau *Edge Universality of β-ensembles*, to appear in Comm. Math. Phys., 2013.

[14] R. Durrett *Probability : Theory and examples*, Fourth edition, Cambridge Univ. Press, 2010.

[15] I. Dumitriu, A. Edelman *Matrix models for beta ensembles.* J. Math. Phys., 43:5830–5847, (2002).

[16] N. Enriquez, L. Ménard *Asymptotic expansion of the expected spectral measure of Wigner matrices*, arXiv.

[17] K. Johansson *On fluctuations of eigenvalues of random Hermitian matrices* Duke Math. Journ. 91, no.1, 151-204 (1998).

[18] R. Killip, M. Stoiciu *Eigenvalue statistics for CMV matrices: From Poisson to clock via random matrix ensembles*, Duke Math. J. 146 (2009), 361–399.

[19] T. Leblé, S. Serfaty *Large Deviation Principle for Empirical Fields of Log and Riesz Gases* arXiv:1502.02970 (2015)

[20] A. Nica, R. Speicher *Lectures on the combinatorics of free probability.* London Mathematical Society Lecture Note Series, 335. Cambridge University Press, Cambridge, 2006.

[21] J.A. Ramírez, B. Rider, B. Virág *Beta ensembles, stochastic Airy spectrum, and a diffusion.* J. Amer. Math. Soc. 24 (2011), 919–944

[22] P. Sosoe, P. Wong, *Local semicircle law in the bulk for Gaussian β-ensemble.* J. Stat. Phys. 148 (2012), no. 2, 204–232.

[23] P. Sosoe, P. Wong, *Convergence of the eigenvalue density for β-Laguerre ensembles on short scales.* Electron. J. Probab. 19 (2014), no. 34, 18 pp.

[24] B. Simon *Trace ideals and their applications*, Mathematical Surveys and Monographs, 120, Second Edition, AMS, 2005.

[25] K.D. Trinh, S. Tomoyuki *The mean spectral measures of random Jacobi matrices related to Gaussian beta ensembles.* arXiv

[26] B. Valkó, B. Virág *Continuum limits of random matrices and the Brownian carousel.* Invent math (2009) 177: 463-508.

MAP 5, UMR CNRS 8145 - Université Paris Descartes, 45 rue des Saints-Pères 75270 Paris cedex 6, France.

E-mail address: `florent.benaych-georges@parisdescartes.fr`

LPMA, Université Paris Diderot, 5 rue Thomas Mann 75013 Paris.

E-mail address: `sandrine.peche@math.univ-paris-diderot.fr`